

COULOMB BLOCKADE FLUCTUATIONS IN DISORDERED QUANTUM DOTS

P. N. WALKER¹, G. MONTAMBAUX² and Y. GEFEN³

¹*Department of Physics and Astronomy, University College London, Gower St., London, England.*

²*Laboratoire de Physique des Solides, associé au CNRS, Université Paris-Sud, 91405 Orsay, France.*

³*Department of Condensed Matter Physics, The Weizmann Institute of Science, 76100 Rehovot, Israel.*

We discuss some recent results on the statistics of the Coulomb Blockade in disordered quantum dots containing spinless interacting fermions using the self-consistent Hartree-Fock approximation. We concentrate on the regime $r_s \gtrsim 1$, with finite dimensionless conductance g . We present significantly different results for the cases of a Coulomb and a nearest-neighbour bare interaction. We show that neglecting rearrangements when the particle number is changed (Koopmans' approximation) can lead to large errors, and show how the spectral statistics of the self-consistent single particle orbitals evolve with the interaction strength.

1 Introduction

The Coulomb Blockade to tunnelling in quantum dots, where the charge in the dot is a good quantum number, provides an experimentally accessible measure of the difference between ground state energies of successive total particle number^{1,2,3,4,5,6}. More specifically, the accessible quantity is the second difference of ground state energies $E_G(N, V_G)$:

$$\Delta_2(N) = E_G(N+1, V_G) - E_G(N, V_G) - E_G(N, V'_G) + E_G(N-1, V'_G), \quad (1)$$

where V_G parametrises the gate voltage environment at each resonant tunnelling point. Within the constant interaction (CI) model the ground state energy is simply the sum of filled single particle energies $\mathcal{E}(n)$ plus $N(N-1)e^2/2C$, where e^2/C is the constant interaction, and $C \propto L^{-1}$ is the geometric capacitance. Taking $V_G = V'_G$, the peak spacing trivially reduces to

$$\Delta_2(N) = \mathcal{E}(N+1) - \mathcal{E}(N) + e^2/C \quad (2)$$

and so, in the diffusive regime, displays Wigner-Dyson (WD) statistics⁷ shifted by e^2/C , up to corrections in one over the dimensionless conductance, g .

Recent experiments on quantum dots^{1,2,3,4} have shown that whilst the mean peak spacings are apparently well described by the CI model, the fluctuations are not described by Wigner-Dyson statistics. It is found that the distribution of Δ_2 is roughly Gaussian^{1,3,4}, with broader non-Gaussian tails seen in Ref. 3. In Refs. 1,2,4 the variance of the fluctuations was found to be considerably larger than that given by the WD distribution. Exact numerical calculations on very small systems¹ also exhibit Gaussian fluctuations on a scale much larger than $\Delta \propto L^{-2}$, and it was suggested that the typical size of the fluctuations scale universally with the average spacing $\langle \Delta_2 \rangle \propto e^2/C$.

In an attempt to bridge the gap between theory⁸ and experiment, Blanter *et al*⁹ evaluated the fluctuations within a Hartree-Fock framework, neglecting effects due to the change in gate voltage V_G to V'_G . To this end they applied the Random Phase Approximation (RPA) to generate the screened interaction in the confined geometry, and assumed that all HF level spacings, except for the Coulomb gap, are described by WD statistics. Implicitly assuming Koopmans' theorem¹⁰ to be valid (which neglects single-particle rearrangements when the particle number distribution is altered), and using wavefunction statistics established for non-interacting electrons in a random potential, they calculated the fluctuations of Δ_2 beyond the CI model. These additional fluctuations were found to be parametrically small (in $1/g$) and proportional to Δ . Hence, the total fluctuations in Δ_2 were found to be proportional to Δ . The disagreement

between experiment and theory therefore remained. The analysis of Ref. 9 is consistent in the limits $g \gg 1$, $r_s \ll 1$. The parameter r_s characterises the relative importance of interactions in the electronic system, and is defined as the mean electron separation in units of the effective Bohr radius. Typical values of $r_s \sim \mathcal{O}(1)$ have been reported in experiment^{1,4}.

To understand why theory and exact numerics are in disagreement, we consider an effective single particle model as in Ref. 9, but generate the single particle orbitals self-consistently using the Hartree-Fock approximation (SCHF). This model allows for the rearrangements when particles are added or removed which was neglected in Ref. 9, and does not require any conjecture about the wavefunction correlations (the two issues are of course intimately linked). We are also able to consider much larger samples than is feasible by exact diagonalisation¹. This approximation has also been seen to be quite good for the calculation of persistent currents in similar systems¹². We show that fluctuations large compared to the single particle level spacing can arise without recourse to varying the sample shape, size or gate-to-dot coupling, supposing these to be *additional* effects. We further demonstrate that approximating the addition spectrum spacings by applying Koopmans' theorem can lead to large errors in the calculation of the spacing statistics. A more detailed presentation of these results can be found elsewhere^{11,13}.

2 The Model

We consider the following tight binding Hamiltonian for spinless fermions

$$H = \sum_i w_i c_i^\dagger c_i - t \sum_{i,\eta} c_{i+\eta}^\dagger c_i + \frac{U}{2} \sum_{ij} M_{ijij} c_i^\dagger c_j^\dagger c_j c_i \quad (3)$$

where i is the site index, η describes the set of nearest neighbours, w_i is the random on site energy in the range $[-W/2, W/2]$, and t the hopping matrix element, henceforth taken as unity. We study separately, both a Coulomb interaction potential,

$$M_{ijij} = 1/|\mathbf{r}_i - \mathbf{r}_j|, \quad (4)$$

and a short range potential,

$$M_{ijij} = \delta_{i,i+\eta}, \quad (5)$$

there is no need to add a constant interaction as it does not affect the physics.

We consider a 2D system with periodic boundary conditions, and choose to define

$$|\mathbf{r}_i - \mathbf{r}_j|^2 \equiv [L_x^2 \sin^2(\pi n_x/L_x) + L_y^2 \sin^2(\pi n_y/L_y)]/\pi^2, \quad (6)$$

where $(n_x, n_y) \equiv \mathbf{r}_i - \mathbf{r}_j$, and L_x, L_y are the sample dimensions.

All distances are measured in units of the lattice spacing a ; the physical parameters are therefore $U = e^2/a$, $t = \hbar^2/2ma^2$. The standard definition for r_s is given, for low filling, by $r_s = U/(t\sqrt{4\pi\nu})$, where $\nu = N/A$ is the filling factor on the tight binding lattice with A sites. The dimensionless conductance g can be approximated, again for low filling, using the Born approximation. We find $g = 96\pi\nu(t/W)^2$, which is valid for $A, N \gg g \gg 1$.

3 The SCHF Approximation

There are a few aspects of this approximation scheme which warrant discussion before presenting the results. Firstly we should stress that this scheme is non-perturbative, but uncontrolled. To ensure that the converged solutions obtained were at least local minima, and hence the RPA excitations stable¹⁴, we diagonalised the total energy curvature matrix¹⁵ in some representative samples. The calculational effort involved is too heavy to do this for all cases.

The issue of which bare potential (4,5) should be used is subtle. If screening is generated externally (by close metallic plates for example), it is reasonable to insert a short-ranged interaction for all interaction strengths. It seems¹³ that this model may then be useful to model the quantum dots of Refs. 5,6. It is not clear however that this form should be taken to represent screening by high-energy particle hole pairs (the RPA approximation). The reason being that the self-consistency process automatically incorporates much of the physics included in the RPA process. In other words, it generates many of the diagrams included in the RPA calculation, and so leads to double counting. This problem does not arise if one applies the Coulomb bare potential. However, we know from basic considerations that the exchange (Fock) interaction should be screened, and the SCHF process is unable to achieve this. In clean systems this can lead to large errors, here however, we have verified that the typical fluctuations of the exchange contribution are smaller than those of the Hartree contribution over the parameter range considered. This suggests that the error made in not screening the exchange term correctly is not overly important. We expect then, that the results for a long ranged interaction should be representative of the exact numerical calculations.

The simplest way to interpret the RPA calculation, is the (average) response of the electron gas to the insertion of a test charge e . Since the excess charge cannot be pushed to infinity in a bounded dot, it is perhaps not surprising that an explicit calculation of the RPA screened interaction shows that a charge e accumulates on the boundary^{9 a}. It was found in Ref. 9 that such *boundary* effects could dominate the interaction induced fluctuations of Δ_2 . In any event, the effect of the boundaries is geometry specific; we remove it by considering a torus geometry.

Since the SCHF approximation attempts to generate static screening by rearranging the effective-single-particle orbitals, there must be a different interpretation between the SCHF approximation with a Coulomb bare potential, and HF with RPA. In the latter, most charge rearrangements are made via the RPA summation, and as such, are separated from the effective-single particles. In our time-independent self-consistent theory, it is not possible to make such a separation, and indeed the job of charge rearrangement is carried purely by the effective-single-particle orbitals. Clearly then, the meaning of the effective-single-particle is not the same in the two cases.

Since the validity of Koopmans' theorem is analysed here, we explain our motivation for investigating an effect that is second order effect in r_s . At sufficiently small r_s , where non-interacting wavefunction correlations can be employed, the effects of rearrangements are negligible. It is often assumed that they are small for the further reason that adding a particle generates only $\mathcal{O}(N)$ corrections to the existing single-particle wavefunctions. We stress however, that when evaluating the ground-state energy, one must sum over N such corrections (as well as N^2 of $\mathcal{O}(N^2)$), and so the cumulative effect can be large. When r_s is sufficiently large as to affect the wavefunction correlations, it is further plausible, that the $\mathcal{O}(r_s^2)$ corrections to the wavefunctions correlate with the unperturbed wavefunctions to generate typical fluctuations of order r_s .

4 Nearest-Neighbour Interactions

In Ref. 11 we show that the distribution of level spacings for both occupied and unoccupied states evolve with r_s from a Wigner-Dyson distribution towards a Poisson-like distribution. This is more pronounced for the unoccupied states, which only have a meaning if Koopmans' approximation is invoked. This tendency is rather weak however, and for $r_s \sim \mathcal{O}(1)$, the distribution is almost indistinguishable from Wigner-Dyson. On the other hand, we found that the self-consistently calculated gap (Δ_2) evolved comparatively rapidly towards a Gaussian

^aWe note that if the dot is supposed to be charge neutral, it may be more consistent if the electron gas were to see an excess background charge of $-e$, plus the *test* electron in question

distribution, and was essentially Gaussian for $r_s \gtrsim 2$. The main difficulty with observing the typical interaction induced fluctuations for $r_s \lesssim 1$ was that they were small compared to Δ . In this regime we confirmed that they scaled like $r_s \Delta$ for fixed g , in agreement with Ref. 9. Indeed, if the wavefunctions remain ergodic, it is clear that this scaling must hold. It was too difficult to identify the disorder scaling with any certainty. In this regime, applying Koopmans' theorem did of course generate errors, but they were found to be relatively small: the mean error scaled like $r_s^2 \Delta$, and the error in the typical fluctuations was too small to analyse with confidence.

When $r_s \gtrsim 1$, we found a dramatic development in the SCHF ground state, and a correspondingly dramatic growth in the error incurred through Koopmans' approximation. We also noted that the system size scaling of the mean Koopmans' error only settled to the $r_s^2 \Delta$ dependence for the larger systems considered ($A \gtrsim 50$ lattice sites); well beyond the current limitations of exact calculations. Whilst the single-particle wavefunctions remained essentially ergodic, incipient charge density modulations (CDMs) led to an unusual, and highly non-universal fluctuation behaviour. In this regime, deviations in $\langle \Delta_2 \rangle$ from the CI model prediction were observed, and identified to be $\mathcal{O}(U/A)$. They were shown to be a direct consequence of the incipient CDMs. These CDMs are analysed in much greater detail in Ref. 13.

In conclusion, we find that for weak interactions, the typical fluctuations $\delta \Delta_2 \sim r_s \Delta$ for fixed $g > 1$, in broad agreement with Blanter *et al*⁹, but not with exact numerics¹. For $r_s \gtrsim 1$, incipient charge density modulations cause a significant change in the statistics of Δ_2 . It is suggested in Ref. 13, that such effects may be relevant to the unusual addition spectra observed in the capacitance experiments of Refs. 5,6.

5 Coulomb Interactions

As for the nearest-neighbour interaction case, the distribution of level spacings for both occupied and unoccupied states were investigated in Ref. 11. They were both found to remain very close to the WD distribution for all interaction strengths considered (up to $r_s \sim 5$). The gap (Δ_2) distribution was found to evolve towards a Gaussian distribution much more rapidly than changes in the above level spacings, but the scale on which this happened depended on the size of the sample.

The mean error generated by Koopmans' approximation was found to tend towards a simple quadratic dependence on r_s as the system size was increased to the maximum size considered of 144 sites and 37 particles. As for the nearest neighbour case, the system size scaling changed around $A \approx 50$ lattice sites. The mean Koopmans' error therefore diverges with system size when compared to both $\langle \Delta_2 \rangle$ and Δ . The Koopmans' approximation error in the typical fluctuations was found to become very large ($\mathcal{O}(\delta \Delta_2)$), no attempt was made to generate an empirical formula to describe it.

The divergent relative error of Koopmans' approximation to the mean charging energy was one of the more striking results of this work. Whilst the CI model worked remarkably well, a supposed improvement on it (Koopmans' theorem about a self-consistent basis) was found to be worse, and moreover, could only be made meaningful for $r_s < \mathcal{O}(1/L)$. This contradicts the expectation that Koopmans' theorem should become a good approximation for $r_s \ll 1$, regardless of the system size. In the CI model this error vanishes identically, while in Blanter *et al*⁹, the terms which generate it were implicitly assumed to vanish.

Turning to the main object of investigation, the SCHF approximation was found to yield typical fluctuations in the charging energy of $\delta \Delta_2 \sim 0.52 \Delta + a \langle \Delta_2 \rangle / \sqrt{g} + \mathcal{O}(r_s^2)$, where a is a constant. A similar result was also found independently by Bonci and Berkovits¹⁶. Whilst this result disagrees with the claim in Ref. 1 that the fluctuations are universal, the system size scaling is consistent. The $\mathcal{O}(r_s^2)$ contribution arises due to interaction induced correlations between the single-particle wavefunctions, and eventually dominates the fluctuations as r_s is

increased. However, the validity of the SCHF approximation becomes more questionable as r_s is increased, and it is not clear to what extent the results are valid in this strong interaction regime. One might expect that the combined fluctuations of the single particle orbitals and the RPA-bosons in the HF with RPA model are similar to the fluctuations generated in the Coulombic SCHF model. Since we do not find agreement with Blanter *et al*⁹, this appears not to be the case. A direct calculation of fluctuations in the polarisability^{17,9} does not account for this discrepancy. It would appear that either screening is not generated in our system (or indeed in the exact calculations on very small samples), or that it is not correct to separate out the RPA-bosons from the fluctuation effects in the usual way.

In summary, we found that Koopmans theorem fails dramatically in the case of a Coulomb bare interaction. We find that the typical interaction induced fluctuations are proportional to $\langle\Delta_2\rangle$ to leading order in r_s , but they are not universal, rather they vanish with the dimensionless conductance like $1/\sqrt{g}$. Other related work has been independently carried out by Levit and Orgad¹⁸, Cohen, Richter and Berkovits¹⁹, and Bonci and Berkovits¹⁶.

Acknowledgments

We acknowledge discussions with Ya. Blanter, A. Finkelstein, S. Levit, C. Marcus, A. Mirlin, D. Orgad, H. Orland, F. von Oppen, F. Piechon, B. Shklovskii and U. Sivan. We acknowledge support from the EU TMR fellowship ERBFMICT961202, the German-Israeli Foundation, the U.S.-Israel Binational-Science Foundation and the Minerva Foundation. Much of the numerical work was performed using IDRIS facilities.

References

1. U. Sivan *et al*, Phys. Rev. Lett. **77**, 1123 (1996).
2. F. Simmel, T. Heinzel and D. A. Wharam, Europhys. Lett. **38**, 123 (1997).
3. S.R. Patel *et al*, Phys. Rev. Lett. **80**, 4522 (1998).
4. F. Simmel *et al*, cond-mat/9901274.
5. R. C. Ashoori *et al*, Phys. Rev. Lett. **68**, 3088 (1992).
6. N. B. Zhitenev *et al*, Phys. Rev. Lett. **79**, 2308 (1997). cond-mat/9703241.
7. M. L. Mehta, Random Matrices 2nd ed., (Academic; New York, 1991).
8. R. Berkovits and B. L. Altshuler Phys. Rev. **B55**, 5297 (1997).
9. Ya. M. Blanter, A. D. Mirlin and B. A. Muzykantskii, Phys. Rev. Lett. **78**, 2449 (1997).
10. T. Koopmans Physica **1**, 104 (1934).
11. P. N. Walker, G. Montambaux and Y. Gefen, cond-mat/9902258.
12. G. Bouzerar and D. Poilblanc, J. Phys. I France **7**, 877 (1997).
13. P. N. Walker, Y. Gefen and G. Montambaux, cond-mat/9902099.
14. D. J. Thouless, Nucl. Phys. **22**, 78 (1961).
15. D. J. Thouless, Nucl. Phys. **21**, 225 (1960).
16. L. Bonci and R. Berkovits cond-mat/9901332.
17. R. Berkovits and B. L. Altshuler Phys. Rev. **B46**, 12 526 (1992).
18. S. Levit and D. Orgad, cond-mat/9901298.
19. A. Cohen, K. Richter and R. Berkovits, cond-mat/9902342.